



Original Research Article

Source-specific Identification and Characterization of Microplastics in Urban Rivers of Surakarta, Indonesia

Darsini¹, Prabang Setyono^{*2}, Mohammad Masykuri³, Bambang Suhardi⁴

¹Department of Environmental Science, School of Postgraduate, Universitas Sebelas Maret, Surakarta, Indonesia

e-mail: darsini.ti@student.uns.ac.id

²Department of Environmental Science, School of Postgraduate, Universitas Sebelas Maret, Surakarta, Indonesia

e-mail: prabangsetyono@staff.uns.ac.id

³Department of Environmental Science, School of Postgraduate, Universitas Sebelas Maret, Surakarta, Indonesia

e-mail: mmasykuri@staff.uns.ac.id

⁴Department of Industrial Engineering, Faculty of Engineering, Universitas Sebelas Maret, Surakarta, Indonesia

e-mail: bambangsuhardi@staff.uns.ac.id

Cite as: Darsini, Setyono, P., Masykuri, M., Suhardi, B., Source-specific Identification and Characterization of Microplastics in Urban Rivers of Surakarta, Indonesia, *J.sustain. dev. energy water environ. syst.*, 14(4), 1140735, 2026, DOI: <https://doi.org/10.13044/j.sdewes.d14.0735>

ABSTRACT

This study investigated the presence and characteristics of microplastics in sediments and surface water from three river sites in Surakarta, Indonesia, influenced by distinct sources of pollution: domestic wastewater, industrial effluents, and agricultural runoff. Samples were collected in triplicate at each site and analysed using stereomicroscopy for morphological classification and Attenuated Total Reflectance–Fourier Transform Infrared Spectroscopy for polymer identification. Microplastic abundances ranged from 1.8 to 3.5 particles/L in surface water and 210 to 385 particles/kg in sediments, with the highest concentrations observed at the domestic site. Polyethylene and polypropylene were the dominant polymers, together accounting for over 65% of all identified particles, while fibers and fragments were the most common morphologies. Variations in polymer type and abundance across sites indicate that different pollution sources shape distinct microplastic profiles. These findings provide critical evidence for targeted interventions at both local and national levels, while also contributing to the global understanding of microplastic pollution in urban rivers. The results highlight the urgent need for integrated management strategies and support policy interventions aligned with Sustainable Development Goals 6 (Clean Water and Sanitation) and Sustainable Development Goals 14 (Life Below Water).

KEYWORDS

Microplastics, Urban rivers, ATR-FTIR, Polymer identification, Southeast Asia.

INTRODUCTION

Microplastics (MPs), defined as plastic particles smaller than 5 mm, have emerged as a pervasive pollutant in aquatic environments worldwide [1]. They can be broadly classified into primary MPs, which are manufactured in small sizes for use in personal care products, industrial abrasives, and pre-production pellets, and secondary MPs, which result from the

* Corresponding author

fragmentation of larger plastic debris through physical, chemical, and biological degradation processes [2]. Major sources of microplastic include domestic activities such as laundry effluents that release synthetic textile fibers [3], industrial discharges containing polymer residues, and the breakdown of mismanaged plastic packaging materials [4]. The physical and chemical characteristics of MPs particularly their size, shape, and density play a critical role in determining their transport, distribution, and environmental fate in freshwater ecosystems [5]. Once ingested by aquatic organisms, MPs can cause physical harm such as gastrointestinal blockage, reduced feeding activity, and mortality, as well as physiological effects including decreased reproductive success and altered growth [6]. Furthermore, MPs can act as vectors for hazardous chemicals and pathogens, increasing the potential ecological and human health risks associated with their presence [7]. Indonesia is widely recognized as one of the most significant contributors of plastic waste to the global ocean. Within this context, the present study advances current knowledge by explicitly identifying and comparing source-specific microplastic discharge pathways in domestic wastewater, industrial effluents, and agricultural runoff within a single urban river system.

Rivers play a pivotal role in transporting microplastics (MPs) from terrestrial environments to marine ecosystems, acting as conduits that integrate inputs from multiple anthropogenic activities along their courses [8]. The primary pathways of MP entry into riverine systems include domestic wastewater discharges, which often contain synthetic fibers from laundry effluents; industrial effluents, which may release polymer fragments, resin pellets, or process-related microplastics; and agricultural runoff [9], which can transport degraded mulch film, irrigation pipe residues, and other synthetic materials [10]. Within river systems, microplastic are distributed between the surface water column and the sediment compartment, with sediment often functioning as a long-term sink due to gravitational settling and hydrodynamic conditions [11]. Factors such as flow velocity, seasonal variation, riparian vegetation, land use, and proximity to pollution sources significantly influence the abundance, distribution, and retention time of MPs in both water and sediments [12].

In Indonesia, research on microplastics (MPs) has predominantly focused on marine and coastal environments, with numerous studies reporting their abundance, polymer types, and potential ecological impacts in seawater, sediments, and biota [13]. Investigations in freshwater systems, particularly urban rivers in Java, remain limited despite their critical role as conduits of plastic waste to the ocean. A recent study provided the first evidence of MPs in the water and sediments of the Surakarta City river basin, highlighting the presence of fibers and fragments predominantly composed of polyethylene (PE) and polypropylene (PP). However, comprehensive assessments linking specific pollution sources—such as domestic wastewater, industrial effluents, and agricultural runoff to variations in MP abundance, morphology, and polymer composition are still scarce in the Indonesian context. Such source-specific studies are essential to develop targeted mitigation strategies, improve waste management practices, and provide baseline data for long-term monitoring of MP pollution in riverine ecosystems [14].

Nonetheless, comprehensive evaluations that connect specific pollution sources, such as domestic wastewater, industrial effluents, and agricultural runoff, to variations in microplastic abundance, morphology, and polymer composition are notably lacking in the Indonesian context. This gap reflects the broader research focus in Indonesia, which has predominantly prioritised marine and coastal environments over freshwater systems [15]. In Southeast Asia, there is a paucity of studies that explicitly correlate the morphology and polymer composition of microplastics with specific pollution sources in riverine environments, despite the region being identified as a global hotspot for plastic waste leakage [16].

Research conducted in Europe and China has demonstrated that microplastic profiles vary systematically according to the predominant anthropogenic activities. For instance, Talbot and Chang demonstrated that land-use patterns and wastewater treatment efficiency significantly influence spatial and temporal variations in microplastic abundance within freshwater systems [17]. Further investigations in China have revealed that industrial activities are primary

contributors to fragment-type microplastics, whereas agricultural regions are frequently associated with plastic films resulting from mulch use [18]. More recent evaluations have underscored that variations in regional waste management practices and socio-economic conditions substantially shape microplastic source signatures in riverine environments [19]. Previous studies have indicated that investigations in freshwater systems, particularly urban rivers in Java, remain limited despite their critical role as conduits of plastic waste to marine environments [10].

Surakarta, located in Central Java, Indonesia, is traversed by several interconnected river systems, including the Bengawan Solo the longest river in Java and its tributaries such as Kali Sroyo, as well as numerous irrigation canals serving surrounding agricultural areas [20]. These waterways receive inputs from diverse anthropogenic activities along their courses. Domestic sources arise from densely populated residential areas that discharge untreated or partially treated household wastewater directly into the rivers, contributing microplastic fibers from laundry effluents and fragments from degraded packaging materials. Industrial sources originate from various manufacturing facilities, including textile, food processing, and plastic production plants, which may release synthetic polymer residues and process-related microplastics into adjacent water bodies [21]. Agricultural sources are associated with runoff carrying degraded mulch films, irrigation hoses, and agrochemical containers, especially during rainfall events [3]. The potential risk is heightened because river water in the region is extensively used for irrigation, aquaculture, and, in some cases, domestic needs, creating direct pathways for microplastic exposure to both aquatic organisms and human populations [2].

Characterizing the morphology and polymer composition of microplastics (MPs) is essential for tracing their sources and assessing potential environmental risks. Variations in shapes such as fibers, often associated with synthetic textiles; fragments, typically derived from the breakdown of larger plastic items; films, commonly originating from packaging materials and agricultural mulch; and pellets, linked to industrial raw materials can provide valuable clues regarding their origin and pathways into aquatic systems [22]. Similarly, identifying the polymer types, including polypropylene (PP), polyethylene (PE), polyethylene terephthalate (PET), polystyrene (PS), and others, is critical for source attribution and for understanding their buoyancy, degradation rates, and potential for adsorbing hazardous chemicals [23]. Among the available analytical techniques, stereomicroscopy is widely employed for initial morphological classification and particle counting, while Fourier-transform infrared spectroscopy (FTIR) has become the standard method for accurate polymer identification due to its ability to detect specific functional groups and differentiate between polymer types [24]. Combining morphological and chemical characterization not only enhances the accuracy of pollution source tracking but also supports ecological risk assessments and the formulation of targeted mitigation strategies [25].

This research provides the first source-specific characterisation of microplastics in Indonesian urban rivers, explicitly linking morphological and polymeric variations to domestic, industrial, and agricultural inputs. This study aims to identify and characterize microplastics present in both sediments and surface water collected from three distinct river sites in Surakarta, each influenced by different dominant pollution sources domestic wastewater, industrial effluents, and agricultural runoff. The research further seeks to analyze the relationships between pollution sources and microplastic characteristics, including their abundance, morphological types (e.g., fibers, fragments, films, pellets), color variations, and polymer compositions as determined through stereomicroscopy and Fourier-transform infrared (FTIR) spectroscopy.

MATERIALS AND METHODS

Sampling was conducted during the dry season in July 2024, under stable hydrological conditions, with no significant rainfall recorded in the 72 hours preceding the sampling. This

period was intentionally selected to minimise short-term hydrological disturbances, such as runoff pulses and sediment resuspension, thereby ensuring comparability among sampling sites. Average ambient temperatures during the sampling period ranged from 26 to 32 °C, and river flow conditions remained relatively stable across all locations. Sampling was performed at three representative river sites in the Surakarta region, each influenced by a distinct dominant pollution source. Sampling was conducted at three representative sites along different river stretches in the Surakarta region. The first site was the Kedung Pedhet River in Mojosongo, Surakarta (7°33'15.5"S 110°50'23.0"E), which is characterized by high population density and dominated by domestic wastewater inputs from residential households. The second site was the Jenes River in Laweyan, Surakarta (7°34'17.2"S 110°47'34.3"E), located in the traditional batik production area, where industrial effluents from batik manufacturing and related small-scale industries constitute the major pollution source. The industrial site is located in Laweyan, a traditional centre of batik production. Batik is a textile industry characterised by wax-resist dyeing processes that predominantly use synthetic dyes, resins, and polymer-based materials. In this study, Sampling Site D (domestic) denotes the domestic wastewater site, Sampling Site I refers to the industrial (batik production) site, and Sampling Site A (agricultural) pertains to the agricultural runoff site. The third site was the Sroyo River in Desa Sroyo, Jaten Subdistrict, Karanganyar (7°32'35.8"S 110°52'38.0"E), which flows through agricultural land and is mainly influenced by agricultural runoff, including residues of mulch films, fertilizers, and pesticides from rice fields and vegetable farms. At each site, both surface water and sediment samples were collected in triplicate during the sampling campaign. The research location map (Figure 1) illustrates the distribution of the three sampling sites, marked as Sampling Site A (agricultural), Sampling Site D (domestic), and Sampling Site I (industry), across the Surakarta region. Triplicate sampling per site was adopted following widely used protocols in microplastic studies, which ensure reproducibility while remaining feasible under logistical constraints. Although larger sample sizes may improve statistical power, triplicates are generally considered adequate for detecting consistent source-specific patterns in microplastic abundance and composition[26].




Figure 1. Map of the sampling area showing Sampling Site D (domestic wastewater), Sampling Site I (industrial/batik production), and Sampling Site A (agricultural runoff)

Materials

The materials used in this study included zinc chloride (ZnCl₂) solution for microplastic extraction from sediments, Whatman No. 41 filter paper with a pore size of 20–25 µm, and distilled water for cleaning all equipment. The instruments and tools comprised a stainless-steel scoop for sediment collection, 300 mL glass bottles for surface water sampling, 500 mL glass beakers, clean polyethylene bags, aluminum foil, a magnetic stirrer, an analytical balance

for precise weighing of sediment samples, a vacuum pump for filtration, standard laboratory glassware, a stereomicroscope with 10–45× magnification for visual identification, and a Fourier Transform Infrared Spectroscopy with Attenuated Total Reflectance (FTIR-ATR) operating over a spectral range of 4000–400 cm.

Methods

Sediment and surface water sampling were conducted at three sites, each representing a different dominant pollution source. At each site, three sediment subsamples were collected from the riverbed at approximately 10 m intervals using a stainless-steel scoop from the top 5 cm layer[26]. The subsamples were wrapped in aluminum foil, sealed in clean polyethylene bags, and transported to the laboratory, where they were air-dried at room temperature, and large debris such as leaves, stones, and shells were manually removed. For microplastic extraction, 20 g of dried sediment from each subsample was placed in a 200 mL glass beaker containing 200 mL of zinc chloride (ZnCl₂) 15% w/v solution. The mixture was stirred for 5 min using a magnetic stirrer, allowed to settle for 30 min, and the supernatant was decanted and filtered through Whatman No. 41 filter paper (pore size 20–25 μm) under vacuum filtration. Surface water sampling involved collecting three 300 mL samples per site at 10 cm depth using pre-cleaned glass bottles, pooling them into a composite sample (900 mL total), and filtering them immediately through the same filter type[27]. All filters were stored in clean Petri dishes until analysis. Visual identification of microplastics was performed under a stereomicroscope at 40× magnification[22], with particles classified by shape (fiber, fragment, film, microbead) and color; to minimize misidentification. Representative particles from each site and sample type were analyzed for polymer composition using Fourier Transform Infrared Spectroscopy with Attenuated Total Reflectance (ATR-FTIR; Bruker model or equivalent), recording spectra over the range of 4000–400 cm⁻¹ at a resolution of 4 cm⁻¹, and matching results against the instrument's spectral library to identify polymer types such as polypropylene (PP), polyethylene (PE), and polyethylene terephthalate (PET) [25]. The total concentrations of microplastics in water and sediment were calculated using the following [question 1](#) and [question 2](#)[28]:

Microplastic concentration in water (particles/L)

$$C_{MP} = \frac{N}{V}$$

Where:

C_{MP} = microplastic concentration (particles/L)
 N = number of particles
 V = volume of water (L)

Microplastic concentration in sediment (particles/kg)

$$C_{MP} = \frac{N}{m}$$

Where:

C_{MP} = microplastic concentration in sediment (particles/kg)
 N = number of particles
 M = mass of sediment (kg)

Due to time and instrument constraints, not all visually identified particles were analysed using FTIR. Instead, a representative subsample of approximately 25–30% of the total particles from each site and matrix (water and sediment) was selected for polymer identification [22], [29]. This approach provides reliable polymer composition profiles while maintaining analytical feasibility.

Quality Assurance and Quality Control

Sediment samples were initially wrapped in aluminium foil before being placed in clean polyethene bags, which functioned solely as secondary external containers. Direct contact between the samples and polyethene was meticulously avoided. Procedural blanks confirmed the absence of polyethene contamination during handling. Rigorous quality assurance protocols were implemented to mitigate potential contamination during sample handling and analysis. Laboratory personnel donned cotton lab coats and nitrile gloves throughout all procedures to minimise the risk of synthetic fibre shedding. All glassware and equipment were meticulously rinsed with pre-filtered distilled water before use. The distilled water employed for cleaning and dilutions was pre-filtered through Whatman GF/F filters (0.7 µm) to eliminate potential microplastic contamination.

Procedural blanks were incorporated by filtering pre-filtered distilled water under identical laboratory conditions and using the same equipment for environmental samples. These blank controls were processed along with each batch of samples to account for potential airborne or procedural contamination. No microplastic particles were detected in the blanks, indicating negligible laboratory contamination during the analyses. These measures align with established international recommendations for minimising laboratory contamination in microplastic studies [22].

Data Analysis

Statistical analyses were conducted to test for significant differences in microplastic abundances among the three sampling sites (domestic, industrial, and agricultural). Normality of the data was first assessed using the Shapiro–Wilk test. For normally distributed data, one-way analysis of variance (ANOVA) was applied, followed by Tukey’s HSD post-hoc test to identify pairwise differences. The non-parametric Kruskal-Wallis test was employed for data that did not meet normality assumptions, with Dunn’s post-hoc test for multiple comparisons. All statistical analyses were performed using R software (version R-4.5.1) with significance set at $p < 0.05$.

RESULTS AND DISCUSSION

[Figure 2](#) shows the variation in microplastic concentrations in surface water across the three sampling sites, reflecting pollution from domestic, industrial, and agricultural sources. The highest concentrations were recorded at the domestic site, likely due to high loads of synthetic fibers from laundry wastewater, detergent residues, and packaging debris entering the river system [27]. Industrial sites exhibited elevated levels of fragments and films, which are commonly associated with manufacturing processes, packaging materials, and polymer-based coatings [26]. Agricultural runoff contributed microplastics primarily in the form of films and colored fragments, consistent with the degradation of polyethylene mulch films and other agroplastic products. Variations in microplastic abundance can also be influenced by hydrodynamic conditions, including flow velocity, seasonal fluctuations, and riparian vegetation cover, which can enhance particle deposition or resuspension [28].




Figure 2. Microplastic concentrations in surface water samples at Sampling Site D (domestic), Sampling Site I (industrial), and Sampling Site A (agricultural)

Figure 3 presents the colour distribution of microplastic particles in surface water, providing insight into their potential sources and environmental weathering processes. Transparent and white particles dominated at the domestic site, consistent with the breakdown of clear packaging films and household plastics. Blue and black fibers were prevalent at the industrial site, potentially originating from synthetic textiles and colored polymer products used in production processes. At the agricultural site, green and brown particles were more common, likely linked to the use of colored mulch films and agricultural twines [26]. The persistence of certain colors may be due to differences in pigment stability, while others fade over time due to UV degradation and biofouling [25].




Figure 3. Microplastic particles categorized by color in water samples

Figure 4 illustrates the concentration of microplastics in sediment samples across the three sampling sites, highlighting sediments as long-term sinks for microplastics. The domestic site exhibited the highest sedimentary concentrations, possibly due to continuous inputs and reduced flushing during low-flow conditions. Industrial sediments contained significant proportions of hard fragments, whereas agricultural sediments had a higher proportion of films. The accumulation of microplastics in sediments is influenced by particle density, size, and shape, as well as local hydrology and sediment composition [26].




Figure 4. Microplastic concentration in sediment samples

Statistical comparisons confirmed that microplastic abundances differed significantly among sites (ANOVA/Kruskal–Wallis, $p < 0.05$). The domestic site exhibited significantly higher concentrations than the industrial and agricultural sites, while no significant difference was detected between the latter. Similar patterns were observed for sediment samples, with the domestic site consistently showing significantly elevated microplastic loads compared to the other locations.

As shown in Figure 2. Microplastic concentrations in surface water samples at Sampling Site D (domestic), Sampling Site I (industrial), and Sampling Site A (agricultural), microplastic concentrations in surface water varied significantly among the three sites. Statistical comparisons confirmed that microplastic abundances differed significantly (ANOVA, $p < 0.05$), with the domestic site exhibiting higher concentrations than the industrial and agricultural sites. Similar patterns were observed in sediments, where Kruskal–Wallis tests indicated significantly greater loads at the domestic site than at other locations. These results are summarised in Table 2. Functional group assignments identified in microplastic particles from sediment samples using ATR-FTIR analysis [9]

Table 1. Results of statistical analysis of differences in microplastic concentrations between locations

Sample type	Test used	df	Test statistic (F / χ^2)	p-value	Post-hoc results ($p < 0.05$)
Surface water (particles/L)	One-way ANOVA	2, N-3	F = 5.87	0.012	Domestic > Industrial, Domestic > Agricultural
Sediment (particles/kg)	Kruskal–Wallis	2	$\chi^2 = 8.34$	0.015	Domestic > Industrial, Domestic > Agricultural

These statistical results are summarised in Table 1. In surface water, ANOVA results indicated that the domestic site exhibited significantly higher abundances than the industrial and agricultural sites ($p < 0.05$). Similarly, Kruskal–Wallis tests for sediment samples showed significantly elevated concentrations at the domestic site, while no significant difference was detected between the industrial and agricultural sites. These findings confirm that the domestic wastewater input is the dominant source of microplastic pollution in Surakarta’s rivers.




Figure 5. Microplastic particles categorized by color in sediment samples

[Figure 5](#) shows the colour distribution of microplastic particles in sediment samples, which largely mirrors the patterns observed in surface water. Transparent and white particles were dominant in domestic sediments, while blue and black were more abundant in industrial sediments. Agricultural sediments showed a greater share of green and brown particles. Over time, biofouling and sediment burial can alter particle appearance, causing originally bright colors to fade or change [26].

ATR-FTIR analysis [Figure 6](#). ATR-FTIR spectra of selected microplastic particles from sediment samples (a) Sampling Site D (domestic), (b) Sampling Site I (industrial), and (c) Sampling Site A (agricultural) identified polyethylene (PE), polypropylene (PP), and polyethylene terephthalate (PET) as the dominant polymers in both surface water and sediment samples ([Figure 6](#)). PE was characterized by strong CH_2 stretching vibrations at 2915 cm^{-1} and 2848 cm^{-1} , CH_2 bending at 1465 cm^{-1} , and CH_2 rocking at 720 cm^{-1} . PP exhibited CH_3 asymmetric stretching near 2950 cm^{-1} and bending modes around 1375 cm^{-1} . PET was identified by $\text{C}=\text{O}$ stretching at 1715 cm^{-1} and aromatic $\text{C}-\text{H}$ bending at 870 cm^{-1} . The polymer distribution patterns suggest a strong link between local human activities and microplastic pollution sources, with domestic inputs contributing more PET from textiles and bottles, industrial sources providing PP from packaging, and agricultural sources introducing PE from mulch films [9].




Figure 6. ATR-FTIR spectra of selected microplastic particles from sediment samples (a) Sampling Site D (domestic), (b) Sampling Site I (industrial), and (c) Sampling Site A (agricultural)

The predominance of fibers at the domestic site can be attributed to untreated laundry effluents, which are known to release large quantities of synthetic microfibers during washing cycles [30]. In Surakarta, the absence of advanced wastewater treatment plants further intensifies microfiber discharge into rivers, as most households discharge greywater directly into nearby waterways. Seasonal variations in laundry frequency, particularly during festive periods, may also contribute to higher microfiber loads. In terms of polymer types, the dominance of PE, PP, and PET reflects local consumption and production patterns. Polyethylene (PE) is widely used in agricultural mulch films and single-use plastic bags, polypropylene (PP) in packaging and household goods, and polyethylene terephthalate (PET) in textiles and bottled beverages, all of which are heavily consumed in Surakarta and throughout Indonesia [31], [32]. These findings indicate that microplastic pollution profiles in Surakarta rivers are not only a function of site-specific waste inputs but also mirror broader socio-economic consumption trends and industrial production activities.

Figure 6 and Table 2 present representative ATR-FTIR spectra of microplastic particles from the three sampling sites, confirming polyethene (PE), polypropylene (PP), and polyethene terephthalate (PET) as the dominant polymers. PE was characterised by strong CH₂ stretching vibrations at 2915 cm⁻¹ and 2848 cm⁻¹, CH₂ bending at 1465 cm⁻¹, and CH₂ rocking at 720 cm⁻¹. PP exhibited CH₃ asymmetric stretching near 2950 cm⁻¹ and bending modes around 1375 cm⁻¹. PET was identified by C=O stretching at 1715 cm⁻¹ and aromatic C-H bending at 870 cm⁻¹. In total, 27% of the visually identified particles were analysed with FTIR, yielding representative spectra that confirmed PE, PP, and PET as the dominant polymers across sites. The polymer distribution patterns suggest a strong link between local human activities and microplastic pollution sources, with domestic inputs contributing more PET from textiles and bottles, industrial sources providing PP from packaging, and agricultural sources introducing PE from mulch films.

Table 2. Functional group assignments identified in microplastic particles from sediment samples using ATR-FTIR analysis [9]

Sampling Site	Wavenumber (cm ⁻¹)	Assignment
---------------	--------------------------------	------------

Site D (Domestic)	717 1027 1372 1459 2850 2915	CH ₂ rock C-O stretch CH ₃ bend CH ₂ bend, CH ₃ bend C-H stretch C-H stretch
Site I (Industrial)	804 839 1162 1367 1454 2833 2915 2946	CH ₂ rock, C-C stretch, C-H stretch CH ₂ rock, C-CH ₃ stretch CH bend, CH ₃ rock, C-C stretch CH ₃ bend CH ₂ bend C-H stretch C-H stretch C-H stretch
Site A (Agricultural)	804 839 1162 1372 1454 2833 2915 2955	CH ₂ rock, C-C stretch, C-H stretch CH ₂ rock, C-CH ₃ stretch CH bend, CH ₃ rock, C-C stretch CH ₃ bend CH ₂ bend C-H stretch C-H stretch C-H stretch

The primary contribution of this study is the quantitative evidence demonstrating that distinct anthropogenic sources produce statistically distinct microplastic fingerprints within a single urban river system. Domestic wastewater was identified as the predominant source of fibres and PET polymers, industrial batik activities were found to contribute disproportionately to PP fragments, and agricultural runoff was primarily linked to PE films. These source-specific profiles were validated through an integrated approach involving morphological, polymeric, and statistical analyses. The detection of microplastics in both water and sediments at all sites underscores their widespread occurrence in urban and peri-urban river systems. Potential ecological impacts include ingestion by aquatic organisms, leading to physical harm, reduced feeding efficiency, and impaired reproduction [25]. Furthermore, microplastics can act as vectors for persistent organic pollutants and heavy metals, exacerbating their toxicity [28]. Effective management strategies should include reducing plastic inputs through wastewater treatment improvements, promoting biodegradable alternatives, and enhancing public awareness campaigns. Long-term monitoring and source-specific mitigation approaches are essential to address microplastic contamination in freshwater ecosystems.

Beyond the quantitative differences observed across sites, the findings also carry significant ecological and socio-economic implications. Ecologically, the high abundance of fibers and fragments in domestic and industrial sites indicates a potential risk for aquatic organisms, as these morphologies are more easily ingested by fish and benthic invertebrates, leading to gastrointestinal blockage, reduced feeding efficiency, and bioaccumulation in food webs [19]. The dominance of polyethylene and polypropylene polymers known for their persistence and buoyancy suggests enhanced mobility in river systems and a higher likelihood of downstream transport to larger rivers and eventually the ocean, thereby contributing to regional-scale plastic leakage [26].

From a socio-economic perspective, Surakarta's rivers provide critical water resources for irrigation, aquaculture, and in some cases, household use. The presence of microplastics in both sediments and surface water implies potential contamination of crops and fish, raising concerns over food safety and public health [33]. This situation could increase economic burdens related

to water treatment and threaten the livelihoods of communities that depend on fisheries and agriculture [30].

The source-specific patterns observed in this study highlight the need for targeted waste management interventions: improving domestic wastewater treatment, enforcing industrial effluent standards, and promoting sustainable agricultural practices. Together, these ecological and socio-economic implications underscore the urgency of integrated policy measures to address microplastic pollution in urban rivers. By linking microplastic characteristics with specific anthropogenic sources, this study provides not only scientific evidence but also actionable insights to guide local authorities and align with broader sustainability frameworks, including SDG 6 (Clean Water and Sanitation) and SDG 14 (Life Below Water).

While prior research conducted in Europe and China has documented microplastic characteristics that vary by source, the current findings reveal a distinct pattern in the prevalence of fibres and PET at domestic locations, indicative of inadequate wastewater treatment infrastructure. In contrast to studies from temperate regions, the greater prevalence of films at agricultural sites underscores the increased use of plastic mulch in tropical agricultural practices. While the study was conducted in Indonesia, its findings are directly relevant to other developing and emerging economies characterised by limited wastewater treatment, small-scale industries, and agriculture with high levels of plastic use. The results indicate that strategies to mitigate microplastic pollution should prioritise filtration of domestic wastewater, implement sector-specific controls on industrial effluents, and manage agricultural plastics. These insights have the potential to inform microplastic policy frameworks beyond Indonesia, particularly in regions such as Southeast Asia, South Asia, Africa, and Latin America.

Policy Implications

The source-specific profiles identified in this study have significant implications for environmental management and policy formulation in Indonesia. Firstly, the predominance of fibres in domestic sites underscores the urgent need to enhance municipal wastewater treatment systems. Installing advanced filtration technologies, such as membrane bioreactors or fine-screening devices, could substantially reduce the discharge of synthetic microfibers from laundry effluents. Pilot projects in major Indonesian cities could serve as models for scaling up such infrastructure nationwide. Secondly, the detection of elevated levels of fragments and films at industrial sites highlights the necessity of enforcing stricter effluent regulations, particularly for the batik industry and other small-scale manufacturers in Surakarta. Developing industry-specific effluent standards that include microplastic parameters, routine monitoring, and non-compliance penalties would help mitigate direct industrial contributions. Incentivising the adoption of closed-loop water recycling systems in batik production could further reduce plastic leakage. Thirdly, the presence of films and colored fragments in agricultural runoff indicates the need for more sustainable management of agro-plastics. Local governments should introduce policies encouraging biodegradable mulch films and establish collection schemes for used irrigation pipes and plastic containers. Extension services could be mobilised to train farmers on best practices in agro-plastic use and disposal, ensuring productivity and environmental sustainability. At a broader scale, integrating microplastic pollution into existing river basin management plans would provide a holistic framework for addressing this issue. Aligning these interventions with national environmental goals and international frameworks such as SDG 6 (Clean Water and Sanitation) and SDG 14 (Life Below Water) would enhance policy coherence and attract international support for implementation.

CONCLUSION

This study provides the first source-specific characterization of microplastics in Surakarta's river systems, demonstrating that domestic, industrial, and agricultural sources each generate

distinct abundance patterns, morphologies, and polymer profiles. Polyethylene and polypropylene dominated across sites, with fibers and fragments as the most common morphologies, indicating both widespread household contributions and industrial inputs. These findings highlight the urgent need for improved wastewater treatment infrastructure, stricter industrial effluent controls, and better management of agricultural plastic use. Beyond Surakarta, the results contribute to the global understanding of microplastic pollution in Southeast Asian urban rivers and support policy interventions aligned with SDG 6 (Clean Water and Sanitation) and SDG 14 (Life Below Water).

ACKNOWLEDGMENT

The authors would like to thank UPT Lab Terpadu, Universitas Sebelas Maret, Surakarta, for providing laboratory facilities and technical assistance during sample preparation and analysis. Appreciation is also extended to the field team members for their support during sampling activities.

NOMENCLATURE

ATR-FTIR	: Attenuated Total Reflectance – Fourier Transform Infrared Spectroscopy
MP	: Microplastic
PE	: Polyethylene
PET	: Polyethylene terphthalate
PP	: Polypropilena
ZnCl ₂	: Zink chloride
SDGs	: Sustainable Development Goals

REFERENCES

- [1] W. Tang, H. Li, L. Fei, B. Wei, T. Zhou, and H. Zhang, "The removal of microplastics from water by coagulation: A comprehensive review," *Sci. Total Environ.*, vol. 851, no. August, p. 158224, 2022, doi: 10.1016/j.scitotenv.2022.158224.
- [2] M. Padervand, E. Lichtfouse, D. Robert, and C. Wang, "Removal of microplastics from the environment . A review," *Environ. Chem. Lett.*, vol. 18, no. 3, pp. 807–828, 2020, doi: 10.1007/s10311-020-00983-1.
- [3] H. S. Auta, C. U. Emenike, and S. H. Fauziah, "Distribution and importance of microplastics in the marine environment : A review of the sources , fate , effects , and potential solutions," *Environ. Int.*, vol. 102, pp. 165–176, 2017, doi: 10.1016/j.envint.2017.02.013.
- [4] R. Geyer, J. R. Jambeck, and K. L. Law, "Production, use, and fate of all plastics ever made - Supplementary Information," *Sci. Adv.*, vol. 3, no. 7, pp. 19–24, 2017, doi: 10.1126/sciadv.1700782.
- [5] Y. Zhang, S. Kang, S. Allen, D. Allen, and T. Gao, "Earth-Science Reviews Atmospheric microplastics : A review on current status and perspectives," *Earth-Science Rev.*, vol. 203, no. September 2019, p. 103118, 2020, doi: 10.1016/j.earscirev.2020.103118.
- [6] S. L. Wright, R. C. Thompson, and T. S. Galloway, "The physical impacts of microplastics on marine organisms : A review," *Environ. Pollut.*, vol. 178, pp. 483–492, 2013, doi: 10.1016/j.envpol.2013.02.031.
- [7] K. D. Cox, G. A. Covernton, H. L. Davies, J. F. Dower, F. Juanes, and S. E. Dudas, "Human Consumption of Microplastics," *Environ. Sci. Technol.*, vol. 53, no. 12, pp. 7068–7074, 2019, doi: 10.1021/acs.est.9b01517.
- [8] L. C. M. Lebreton, J. Van Der Zwet, J. Damsteeg, B. Slat, A. Andrady, and J. Reisser, "River plastic emissions to the world ' s oceans," *Nat. Commun.*, vol. 8, pp. 1–10, 2017,

- doi: 10.1038/ncomms15611.
- [9] A. Dilshad, M. Taneez, F. Younas, and A. Jabeen, "Microplastic pollution in the surface water and sediments from Kallar Kahar wetland , Pakistan : occurrence , distribution , and characterization by ATR - FTIR," *Environ. Monit. Assess.*, 2022, doi: 10.1007/s10661-022-10171-z.
- [10] A. Ismanto, T. Hadibarata, D. Nugroho, M. Zainuri, and A. Mehmood, "First evidence of microplastics in the water and sediment of Surakarta city river basin , Indonesia," *Mar. Pollut. Bull.*, vol. 196, no. September, p. 115677, 2023, doi: 10.1016/j.marpolbul.2023.115677.
- [11] S. Klein, E. Worch, and T. P. Knepper, "Microplastics in the Rhine-Main area in Germany : Occurrence , spatial distribution and sorption of organic contaminants," *Environ. Sci. Technol.*, vol. 49, no. 0, pp. 2–3, 2015.
- [12] Z. Li et al., "Association of ambient air pollutants and birth weight in Ningbo ," *Environ. Pollut.*, vol. 249, pp. 629–637, 2019, doi: 10.1016/j.envpol.2019.03.076.
- [13] J. Beyer, H. C. Trannum, T. Bakke, P. V Hodson, and T. K. Collier, "Environmental effects of the Deepwater Horizon oil spill : A review," *MPB*, vol. 110, no. 1, pp. 28–51, 2016, doi: 10.1016/j.marpolbul.2016.06.027.
- [14] A. Muñoz-vera, J. Matías, P. Castejón, and G. García, "Patterns of trace element bioaccumulation in jelly fi sh *Rhizostoma pulmo* (Cnidaria , Scyphozoa) in a Mediterranean coastal lagoon from SE Spain," *MPB*, vol. 110, no. 1, pp. 143–154, 2016, doi: 10.1016/j.marpolbul.2016.06.069.
- [15] A. A. Koelmans, N. H. Mohamed Nor, E. Hermsen, M. Kooi, S. M. Mintenig, and J. De France, "Microplastics in freshwaters and drinking water: Critical review and assessment of data quality," *Water Res.*, vol. 155, pp. 410–422, 2019, doi: 10.1016/j.watres.2019.02.054.
- [16] A. Shen et al., "Current status of microplastic pollution in China's aquatic environment and its interactions with metal pollutants on aquatic organisms," *Water Res. X*, vol. 28, p. 100392, 2025, doi: 10.1016/j.wroa.2025.100392.
- [17] H. Zhao et al., "Pollution status of microplastics in the freshwater environment of China: a mini review," *Water Emerg. Contam. Nanoplastics*, vol. 1, no. 1, 2022, doi: 10.20517/wecn.2021.05.
- [18] R. Talbot and H. Chang, "Microplastics in freshwater: A global review of factors affecting spatial and temporal variations," *Environ. Pollut.*, vol. 292, no. April 2021, p. 118393, 2022, doi: 10.1016/j.envpol.2021.118393.
- [19] F. M. Windsor, I. Durance, A. A. Horton, R. C. Thompson, C. R. Tyler, and S. J. Ormerod, "A catchment-scale perspective of plastic pollution," *Glob. Chang. Biol.*, vol. 25, no. 4, pp. 1207–1221, 2019, doi: 10.1111/gcb.14572.
- [20] A. Bakir et al., "Occurrence and abundance of meso and microplastics in sediment , surface waters , and marine biota from the South Pacific region," *Mar. Pollut. Bull.*, vol. 160, no. August, p. 111572, 2020, doi: 10.1016/j.marpolbul.2020.111572.
- [21] M. Firdaus, Y. Trihadiningrum, and P. Lestari, "Microplastic pollution in the sediment of Jagir Estuary , Surabaya City , Indonesia," *Mar. Pollut. Bull.*, vol. 150, no. September 2019, p. 110790, 2020, doi: 10.1016/j.marpolbul.2019.110790.
- [22] V. Hidalgo-ruz, L. Gutow, R. C. Thompson, and M. Thiel, "Microplastics in the Marine Environment: A Review of the Methods Used for Identification and Quantification," *Environ. Sci. Technol.*, vol. 46, pp. 3060–3075, 2012, doi: 10.1021/es2031505.
- [23] G. Lamichhane et al., "Microplastics in environment: global concern, challenges, and controlling measures," *Int. J. Environ. Sci. Technol.*, vol. 20, no. 4, pp. 4673–4694, 2023, doi: 10.1007/s13762-022-04261-1.
- [24] M. Gu, M. Kuczera, and S. Mintenig, "Focal plane array detector-based micro-Fourier-transform infrared imaging for the analysis of microplastics in environmental samples," *Environ. Chem.*, vol. 12, pp. 563–581, 2015.

- [25] P. Costa, A. C. Duarte, T. Rocha-santos, and J. C. Prata, "Trends in Analytical Chemistry Methods for sampling and detection of microplastics in water and sediment : A critical review Density separation," *Trends Anal. Chem.*, vol. 110, pp. 150–159, 2019, doi: 10.1016/j.trac.2018.10.029.
- [26] R. R. Hurley, A. L. Lusher, M. Olsen, and L. Nizzetto, "Validation of a Method for Extracting Microplastics from Complex, Organic-Rich, Environmental Matrices," *Environ. Sci. Technol.*, vol. 52, pp. 7409–7417, 2018, doi: 10.1021/acs.est.8b01517.
- [27] C. M. Free, O. P. Jensen, S. A. Mason, M. Eriksen, and N. J. Williamson, "High-levels of microplastic pollution in a large , remote , mountain lake," *Mar. Pollut. Bull.*, vol. 85, no. 1, pp. 156–163, 2014, doi: 10.1016/j.marpolbul.2014.06.001.
- [28] D. E. Eppehimer *et al.*, "Impacts of baseflow and flooding on microplastic pollution in an effluent-dependent arid land river in the USA," *Environ. Sci. Pollut. Res.*, vol. 28, pp. 45375–45389, 2021.
- [29] J. C. Prata, J. P. da Costa, A. C. Duarte, and T. Rocha-Santos, "Methods for sampling and detection of microplastics in water and sediment: A critical review," *TrAC - Trends Anal. Chem.*, vol. 110, pp. 150–159, 2019, doi: 10.1016/j.trac.2018.10.029.
- [30] I. E. Napper and R. C. Thompson, "Release of synthetic microplastic plastic fibres from domestic washing machines: Effects of fabric type and washing conditions," *Mar. Pollut. Bull.*, vol. 112, no. 1–2, pp. 39–45, 2016, doi: 10.1016/j.marpolbul.2016.09.025.
- [31] J. R. Jambeck *et al.*, "Plastic waste inputs from land into the ocean," *Ciencia*, vol. 347, no. 6223, pp. 768–771, 2015. <https://doi.org/10.1126/science.1260352>
- [32] M. R. Cordova, T. A. Hadi, and B. Prayudha, "Occurrence and abundance of microplastics in coral reef sediment : a case study in Sekotong , Lombok-Indonesia," *Adv. Environ. Sci. Bioflux*, vol. 10, no. 1, pp. 23–29, 2018, doi: 10.5281/zenodo.1297719.
- [33] C. M. Rochman *et al.*, "Anthropogenic debris in seafood : Plastic debris and fibers from textiles in fish and bivalves sold for human consumption," *Nat. Publ. Gr.*, no. September, pp. 1–10, 2015, doi: 10.1038/srep14340.